arXiv:1604.00142v1 [math.DG] 1 Apr 2016

ASYMPTOTIC BEHAVIOR OF POSITIVELY CURVED STEADY RICCI SOLITONS, II

YUXING DENG AND XIAOHUA ZHU*

Abstract. In this note, we show that any n-dimensional κ-noncollapsed steady Kähler-Ricci soliton with nonnegative bisectional curvature must be flat. The result is an improvement to our former work in [7].

1. Introduction

In [7], we prove

Theorem 1.1. *There is no any κ-noncollapsed steady Kähler-Ricci soliton with nonnegative sectional curvature and positive Ricci curvature.*

In this note, we improve Theorem 1.1 in sense of nonnegative bisectional curvature. Namely, we have

Theorem 1.2. *There is no any κ-noncollapsed steady Kähler-Ricci soliton with nonnegative bisectional curvature and positive Ricci curvature.*

As an application of Theorem 1.1, we get the following rigidity result for the steady Kähler-Ricci solitons and Kähler-Ricci flow.

Theorem 1.3. *Any κ-noncollapsed steady Kähler-Ricci soliton (M, g, f) with nonnegative bisectional curvature must be flat. More generally, any κ-noncollapsed noncompact and eternal Kähler-Ricci flow with nonnegative bisectional curvature and uniformly bounded curvature must be a flat flow.*

In the proof of Theorem 1.1, one main step is to use the blow-down argument to analysis the structure of limit Kähler-Ricci flow of a sequence of rescaled Kähler-Ricci flows. The nonnegative sectional curvature condition is used to guarantee the existence of lines on the limit flow by using Toponogov comparison Theorem (cf. [9]) and consequently the limit flow can

2000 *Mathematics Subject Classification.* Primary: 53C25; Secondary: 53C55, 58J05.
Key words and phrases. Ricci flow, Ricci soliton, κ-solution.
* Partially supported by the NSFC Grants 11271022 and 11331001.

be split off a line by Cheeger-Gromoll splitting theorem [4]. The splitting property is crucial in our proving the curvature decay:

$$R(p,t) \to 0, \ as \ t \to -\infty. \tag{1.1}$$

Here $R(p,t)$ is a scalar curvature of corresponding Ricci flow of steady Kähler-Ricci soliton (M,g,f).

The new ingredient at present is to prove a local splitting result for κ-noncollapsed steady Kähler-Ricci solitons with nonnegative bisectional curvature (cf. Lemma 3.1). Then we can generalize the argument in the proof of Theorem 1.1 to Theorem 1.2.

Theorem 1.3 can be regarded as a generalization of Ni's rigidity theorem for ancient solution of Kähler-Ricci flow with maximal volume growth [10]. Ni's result is a complex version of Perelman's Theorem for ancient solution of Ricci flow in [11], Section 11.4.

Acknowledgements. The work was done partially when the second author is visiting at the Mathematical Sciences Research Institute at Berkeley during the spring 2016 semester. The author would like to thank her hospitality and the financial supports, National Sciences Foundation under Grant No. DMS-1440140, and Simons Foundation.

2. Primary results in [7]

In this section, we recall some results proved in [7], which will be used to complete the proof of Theorem 1.2 in the next section.

Let (M,g,f) be a complete κ-noncollapsed steady Kähler-Ricci soltion with nonnegative bisectional curvature. It is proved that there exists a quilibrium point o of M such that $\nabla f(o)=0$ in [6]. Let ϕ_t be a family of biholomorphisms generated by $-\nabla f$. Let $g(t)=\phi_t^*(g)$. Then $g(t)$ satisfies the Ricci flow

$$\frac{\partial g}{\partial t} = -2\mathrm{Ric}(g), \ t \in (-\infty,\infty). \tag{2.1}$$

Let $p_i \in M$ be any sequence with $\mathrm{dist}(o,p_i) \to \infty$ as $i \to \infty$. We consider a sequence of rescaled flows $(M, R(p_i)g(R^{-1}(p_i)t), p_i)$. By using the compactness theorem, Theorem 3.3 in [7], we prove the following convergence result.

Theorem 2.1. *$(M, R(p_i)g(R^{-1}(p_i)t), p_i)$ converges subsequently to a pseudo κ-solution $(M_\infty, g_\infty(t))$ $(t \in (-\infty,\infty))$ of Kähler-Ricci flow in the Cheeger-Gromov topology.*

Theorem 2.1 is in fact of a part of Theorem 1.5 in [7]. Since (M,g,f) is weakened to have nonnegative bisectional curvature condition, we could not

get the splitting property of limit $(M_\infty, g_\infty(t))$ as in Theorem 1.5 under nonnegative sectional curvature condition. Here a pseudo κ-solution of Kähler-Ricci flow means that a non-flat Kähler-Ricci flow with κ-noncollapsed condition satisfies the Harnack inequality,

$$\frac{\partial R}{\partial t} + \nabla_i R V^i + \nabla_{\bar{i}} R V^{\bar{i}} + R_{i\bar{j}} V^i V^{\bar{j}} \geq 0, \ \forall \ V \in T^{(1,0)}M. \tag{2.2}$$

For e a steady Kähler-Ricci soliton (M, g, J) with positive Ricci curvature, which admits an equilibrium point, Bryant in [1] proves that there exist global holomorphic coordinates (Poincaré coordinates) $z : M \to \mathbb{C}^n$ which linearize $Z = \frac{\nabla f - \sqrt{-1}J\nabla f}{2}$ such that

$$Z = \sum_{i=1}^{n} h_i z_i \frac{\partial}{\partial z_i}, \tag{2.3}$$

where $h_1, \cdots, h_n$ are positive constants. As in Corollary 5.2 in [7], we choose points $p_k = (k, 0, \cdots, 0, 0 \cdots, 0) \in M$. By computation of lengths of integral curves of $J\nabla f$ starting from p_k, we have

Lemma 2.2. *Suppose that (M, g, f) is an n-dimensional κ-noncollapsed steady Kähler-Ricci soliton with nonnegative bisectional curvature and positive Ricci curvature. Then there exists a positive constant C such that $R(p_k) > C$, where C is independent of p_k.*

Lemma 2.2 is in fact from Lemma 5.4 in [7] while the nonnegative sectional curvature condition is replaced by nonnegative bisectional curvature condition. This can be done since we only use the convergence result, Theorem 2.1 in the proof Lemma 5.4.

3. Proof of Theorem 1.2

The following proposition is a key lemma in the proof of Theorem 1.2.

Lemma 3.1. *Let (M, g, f) be a κ-noncollapsed steady Kähler-Ricci soltion with nonnegative bisectional curvature and positive Ricci curvature. Then, for any $p \in M \setminus \{o\}$, at least one of the following two properties holds:*

i) $R(p, t) \to 0$, as $t \to -\infty$.

ii) For any $\tau_i \to -\infty$, $(M, R(p, \tau_i)g(R^{-1}(p, \tau_i)t), \phi_{\tau_i}(p))$ converges subsequently to a flow of κ-noncollapsed steady Kähler-Ricci soltions $(M_\infty, g_\infty(t)$, $p_\infty)$ with nonnegative bisectional curvature which splits locally off a complex line.

Proof. First we note

$$|\nabla f|^2 + R = R_{\max} = \sup_{x \in M} R(x). \tag{3.1}$$

and $R(p) \geq 0$. Then by the uniqueness of quilibrium points, we have

$$0 < R(p,t) < R_{\max}.$$

Thus by relation

$$\frac{\partial R(p,t)}{\partial t} = 2\mathrm{Ric}(\nabla f, \nabla f)(\phi_t(p)) > 0, \tag{3.2}$$

we see that $\lim_{t\to-\infty} R(p,t)$ exists. Now we suppose that the property i) in Lemma 3.1 does not hold. Then there exists a point $p \in M$ such that

$$\lim_{t\to-\infty} R(p,t) = C_0 > 0. \tag{3.3}$$

Since o is an unique quilibrium point, by (3.2), we have $C_0 < R_{\max}$.

Consider any sequence $(M, g_i(t), p_{\tau_i})$, where $g_i(t) = R(p,\tau_i)g(R^{-1}(p,\tau_i)t)$ and $p_{\tau_i} = \phi_\tau(p)$ with $\tau_i \to -\infty$. Then $(M, g_i(t))$ is κ-noncollapsed and curvature of $(M, g_i(t))$ is uniformly bounded. By Hamilton compactness theorem [8], $(M, g_i(t), p_{\tau_i})$ converges subsequently to a pseudo κ-solution $(M_\infty, g_\infty(t), p_\infty)$, where $t \in (-\infty, \infty)$. Moreover, by (3.3), we have

$$R_\infty(p_\infty, t) = \lim_{\tau_i\to-\infty} \frac{R(p_{\tau_i}, R^{-1}(p,\tau_i)t)}{R(p,\tau_i)} = 1, \ \forall\ t \in (-\infty, +\infty). \tag{3.4}$$

and consequently,

$$\frac{\partial}{\partial t} R_\infty(p_\infty, t) \equiv 0. \tag{3.5}$$

Since $(M_\infty, g_\infty(t); p_\infty)$ is not flat by (3.4), we may assume that $(M_\infty, g_\infty(t))$ has positive Ricci curvature by Cao's dimension reduction theorem [3]. By the Harnack inequality (2.2) together with condition (3.5), following the argument in the proof of Theorem 4.1 in [2], we can further prove that $(M_\infty, g_\infty(t), p_\infty)$ is in fact a steady Kähler-Ricci soliton, which is κ-noncollapsed and has nonnegative bisectional curvature and positive Ricci curvature (also see Proposition 2.2, [5]). More precisely, there is a smooth vector field V on M_∞ such that

$$R^{(\infty)}_{ij} = \frac{1}{2}(\nabla^{(\infty)}_i V_j + \nabla^{(\infty)}_j V_i) \tag{3.6}$$

and

$$\nabla^{(\infty)}_i V_j = \nabla^{(\infty)}_j V_i. \tag{3.7}$$

Thus

$$\frac{\partial R^{(\infty)}(p_\infty, t)}{\partial t}|_{t=0} = 2\mathrm{Ric}^{(\infty)}(V, V).$$

By (3.5), it follows $V(p_\infty) = 0$.

Let $X_{(i)} = R^{-1}(p,\tau_i)\nabla f$. Then

$$|\nabla^i X_{(i)}|_{\hat{g}_i} \leq C_0^{-1}|\mathrm{Ric}^{(i)}|_{\hat{g}_i} \leq C,$$

where $\hat{g}_i = R(p,\tau_i)g$ and $\mathrm{Ric}^{(i)}$ is Ricci curvature of $\hat{g}_i$. By Shi's higher order estimate [12] and the soliton equation, we also get

$$|(\nabla^i)^m X_{(i)}|_{\hat{g}_i} \le C(n)|(\nabla^i)^{m-1}\mathrm{Ric}^{(i)}|_{\hat{g}_i} \le C(m).$$

Thus by taking a subsequence, we may assume that

$$X_{(i)} \to X \ as \ i \to \infty.$$

Since

$$\nabla^{(i)}_j X_{(i)k} = \nabla^{(i)}_k X_{(i)j} = R^{(i)}_{jk},$$

we get

$$\nabla^{(\infty)}_j X_k = \nabla^{(\infty)}_k X_j = R^{(\infty)}_{jk}. \tag{3.8}$$

Moreover, by (3.1),

$$|X|_{g_\infty}(p_\infty) = \lim_{i\to\infty} |X_{(i)}|_{g_i}(p_i) = \lim_{i\to\infty}\sqrt{\frac{R_{\max}}{R(p,t_i)} - 1}.$$

Hence

$$|X|_{g_\infty}(p_\infty) = \sqrt{C_0^{-1}R_{\max} - 1} > 0. \tag{3.9}$$

Let $W = V - X$. Then by (3.6), (3.7) and (3.8), we have

$$\nabla^{(\infty)} W = \nabla^{(\infty)}(V - X) \equiv 0. \tag{3.10}$$

On the other hand, $V(p_\infty) = 0$. By (3.9), we see that W is nonzero everywhere. Thus the Kähler manifold $(M_\infty, g_\infty(0))$ splits locally off a line along W. Note that $J_\infty W(p_\infty) \neq 0$ and

$$\nabla^{(\infty)}(J_\infty W) = 0,$$

where J_∞ is the complex structure of M_∞. Hence $(M_\infty, g_\infty(0))$ also splits locally off a line along $J_\infty W$. As a conseqence, the steady soliton $(M_\infty, g_\infty(0))$ splits locally off a complex line. Therefore, the property ii) in Lemma 3.1 holds. The lemma is proved. □

Remark 3.2. *Lemma 3.1 is still true for the steady Kähler-Ricci soliton with nonnegative bisectional curvature if the κ-noncollapsing condition is replaced by the existence of uniform injective radius of soliton.*

Corollary 3.3. *Let (M, g, f) be a 2-dimensional κ-noncollapsed steady Kähler-Ricci soliton with nonnegative bisectional curvature and positive Ricci curvature. Then for any $p \neq o$,*

$$R(p,t) \to 0, \ \text{as } t \to -\infty. \tag{3.11}$$

Proof. We prove the corollary by contradiction. Then by Lemma 3.1 there exist a point $p \in M$ and a sequence of $t_i \to -\infty$ such that

$$\lim_{t\to-\infty} R(p,t_i) = C > 0 \tag{3.12}$$

and rescaled $(M, R(p,t_i)g(R^{-1}(p,t_i)t), \phi_{t_i}(p))$ converges subsequently to a flow of 2-dimensional complete κ-noncollapsed, steady Kähler-Ricci solitons $(M_\infty, g_\infty(t), p_\infty)$ $(t \in (-\infty,\infty))$ which splits locally off a complex line. Moreover

$$R_\infty(p_\infty, t) = 1, \ for\ t \in (-\infty, +\infty). \tag{3.13}$$

Let $(\widetilde{M}_\infty, \widetilde{g}(0))$ be the universal covering of $(M_\infty, g(0))$. Then it is easy to see that $(\widetilde{M}_\infty, \widetilde{g}(0))$ still satisfies the κ-noncollapsing condition. Moveover, the parallel vector field W in Lemma 3.1 can be lifted on $(\widetilde{M}_\infty, \widetilde{g}(0))$ and this vector field generates a trivial honolomy group of $(\widetilde{M}_\infty, \widetilde{g}(0))$. Thus by Wu's de Rham decomposition Theorem [13], the steady soliton $(\widetilde{M}_\infty, \widetilde{g}(0))$ splits off a complex line. Hence the corresponding steady solitons flow $(\widetilde{M}_\infty, \widetilde{g}(t))$ splits off a complex line, and so it splits out a real 2-dimensional flow $(N, g_N(t))$ of complete κ-noncollapsed steady Ricci solitons with positive curvature. On the other hand, by Lemma 4.4 in [7], any complete pseudo-κ-solution on a surface is a shrinking flow of round spheres and so $(N, g_N(t))$ does. In particular, $(N, g_N(0))$ is compact. But this is impossible since any compact gradient steady Ricci soliton should be flat. The corollary is proved.

□

Choosing $p \neq o \in M$ and a sequence of $t_k \to -\infty$. Let $p_k = \phi_{t_k}$. Then by Corollary 3.3, $R(p,t_k) \to 0$ as $t_k \to -\infty$. Moreover, one can check $p_k = (e^{-t_k h_1}, ..., 0)$ under Poincaré coordinates. Thus applying Lemma 2.2 to 2-dimensional steady Kähler-Ricci soliton together with Cao's dimension reduction theorem, we prove

Proposition 3.4. *Let (M,g) be a 2-dimensional κ-noncollapsed steady Kähler-Ricci soliton with nonnegative bisectional curvature. Then (M,g) is flat.*

Now we complete the proof of Theorem 1.2.

Proof of Theorem 1.2. By Proposition 3.4, we can use the induction argument as in the proof of Theorem 1.3, [7]. Suppose that there is no k-dimensional κ-noncollapsed steady Kähler-Ricci soliton with nonnegative bisectional curvature and positive Ricci curvature for all $k < n$. We claim

Claim 3.5. *Under the induction hypothesis, for any fixed $p \in M \setminus \{o\}$, $R(p,-t) \to 0$ as $t \to \infty$.*

The proof of Claim 3.5 is similar to one of Corollary 3.3. In fact, if the claim is not true, then by Lemma 3.1 there exist a point $p \in M$ and a sequence of $t_i \to -\infty$ such that

$$\lim_{t\to-\infty} R(p,t_i) = C > 0 \tag{3.14}$$

and rescaled $(M, R(p,t_i)g(R^{-1}(p,t_i)t), \phi_{t_i}(p))$ converges subsequently to a flow $(M_\infty, g_\infty(t), p_\infty)$ $(t \in (-\infty,\infty))$ of complete κ-noncollapsed, steady Kähler-Ricci solitons which splits locally off a complex line. Moreover

$$R_\infty(p_\infty, t) = 1, \ for\ t \in (-\infty, +\infty). \tag{3.15}$$

Now we can consider the universal covering $(\widetilde{M}_\infty, \widetilde{g}(0))$ of $(M_\infty, g(0))$. As in Corollary 3.3, $(\widetilde{M}_\infty, \widetilde{g}(0))$ can be split out an $n-1$-dimensional complete κ-noncollapsed steady Kähler-Ricci soliton (N, g_N) with nonnegative bisectional curvature. Note that (N, g_N) is not flat by (3.15). Thus by Cao's dimension reduction theorem, we may assume that (N, g_N) has positive Ricci curvature. On the other hand, by the induction assumpation, (N, g_N) should be flat. This is a contradiction! The claim is proved.

By Claim 3.5, we can choose p_k as in the proof of Proposition 3.4 to apply Lemma 2.2 to finish the proof of Theorem 1.2 .

□

Theorem 1.3 is an application of Theorem 1.2 by using an argument as in proof of Theorem 1.4 in [7].

References

[1] Bryant, R. L., *Gradient Kähler Ricci solitons*, Géomtrié différentielle, physique mathématique, mathématiques et société. I. Astérisque, No. 321 (2008), 51-97.

[2] Cao, H.D., *Limits of solutions to the Kähler-Ricci flow*, J. Diff. Geom., **45** (1997), 257-272.

[3] Cao, H.D., *On dimension reduction in the Kähler-Ricci flow*, Comm. Anal. Geom., **12**, No. 1, (2004), 305-320.

[4] Cheeger, J. and Gromoll, D., *The splitting theorem for manifolds of nonnegative Ricci curvature*, J. Diff. Geom., **6** (1972), 119-128.

[5] Chau, A. and Tam, L-F., *Non-negatively curved Kähler manifolds with average quadratic curvature decay*, Comm. Anal. Geom. **15** (2007), no. 1, 121-146.

[6] Deng, Y.X. and Zhu, X.H., *Complete non-compact gradient Ricci solitons with nonnegative Ricci curvature*, Math. Z., **279** (2015), no. 1-2, 211-226.

[7] Deng, Y.X. and Zhu, X.H., *Asymptotic behavior of positively curved steady Ricci solitons*, arXiv:math/1507.04802.

[8] Hamilton, R.S., *Formation of singularities in the Ricci flow*, Surveys in Diff. Geom., **2** (1995), 7-136.

[9] Morgan, J. and Tian, G., *Ricci flow and the Poincaré conjecture*, Clay Math. Mono., 3. Amer. Math. Soc., Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2007.

[10] Ni, L., *Ancient solutions to Kähler-Ricci flow*, Math. Res. Lett., 12 (2005), 633-654.

[11] Perelman, G., *The entropy formula for the Ricci flow and its geometric applications*, arXiv:math/0211159.

[12] Shi, W.X., *Ricci deformation of the metric on complete noncompact Riemannian manifolds*, J. Diff. Geom., **30** (1989), 223-301.

[13] Wu, H., *On the de Rham decomposition Theorem*, Illinois J. Math., **8** (1964), 291-311.

Yuxing Deng, School of Mathematical Sciences, Beijing Normal University, Beijing, 100875, China, dengyuxing@mail.bnu.edu.cn

Xiaohua Zhu, School of Mathematical Sciences and BICMR, Peking University, Beijing, 100871, China, xhzhu@math.pku.edu.cn